

**THE FOLLOWING ARE THE ENGLISH TRANSLATION
OF ANNEXES TO THE INTERNATIONAL PRELIMINARY
EXAMINATION REPORT (ARTICLE 34):**

Amended Sheets (Pages 51-54)

AS ENCLOSED TO IPRP

We claim:

- 5 1. A catalytically active composition comprising an active component having the formula



10 wherein Y = Au or Rh,
 and wherein the indices a, c and d indicate the mass ratios of the respective elements and $0.1 \leq a \leq 3$, $0.1 \leq c \leq 3$ and $0 \leq d \leq 1$,
 on silicon carbide or steatite as carrier.

- 15 2. A catalytically active composition comprising an active component having the formula




20 wherein the indices a, b, c indicate the mass ratios of the respective elements
 and $0.1 \leq a \leq 3$, $0 \leq b \leq 3$ and $0.1 \leq c \leq 3$,
 on silicon carbide or steatite as carrier.

- 25 3. A catalytically active composition comprising an active component having the formula



30 wherein a and c indicate the mass ratios of the respective elements and $0.1 \leq a \leq 3$ and $0.1 \leq c \leq 3$,
 on silicon carbide or steatite as carrier.

- 35 4. A catalytically active composition comprising an active component having the formula



wherein Z = Ag or Pt,

- 52 -

and wherein the indices a, b, c and e indicate the mass ratios of the respective elements and $0.1 \leq a \leq 3$, $0 \leq b \leq 3$, $0.1 \leq c \leq 3$ and $0 \leq e \leq 1$, on silicon carbide or steatite as carrier.

- 5 5. A catalytically active composition comprising an active component having the formula



10 wherein the indices a, c and e indicate the mass ratios of the respective elements and $0.1 \leq a \leq 3$, $0.1 \leq c \leq 3$ and $0 \leq e \leq 1$, on silicon carbide or steatite as carrier.

- 15 6. A catalytically active composition comprising an active component of the formula:

- 20 - $\text{Pd}_{0.5-1.0}\text{Rh}_{0.5-1.25}\text{Bi}_{1.25-1.75}\text{Ag}_{0.05-0.15}$
 - $\text{Pd}_{0.5-1.0}\text{Rh}_{1.0-1.5}\text{Bi}_{0.75-1.25}\text{Pt}_{0.01-0.1}$
 - $\text{Pd}_{0.25-0.5}\text{Rh}_{1.75-2.5}\text{Bi}_{0.25-0.5}\text{Co}_{0.01-0.1}$
 - $\text{Pd}_{0.5-1.25}\text{Rh}_{0.5-1.25}\text{Bi}_{0.75-1.5}\text{Cr}_{0.01-0.1}$
 - $\text{Pd}_{1.0-1.75}\text{Rh}_{0.25-0.75}\text{Bi}_{0.75-1.5}\text{Pt}_{0.0-0.15}\text{Co}_{0.01-0.1}$
 - $\text{Pd}_{1.0-1.75}\text{Rh}_{0.25-0.75}\text{Bi}_{0.75-1.5}\text{Pt}_{0.05-0.15}$
 - $\text{Pd}_{0.5-1.0}\text{Rh}_{1.0-1.75}\text{Bi}_{0.5-1.25}\text{Ag}_{0.03-0.15}\text{Ca}_{0.02-0.1}$
 - $\text{Pd}_{0.4-1.0}\text{Rh}_{1.0-1.75}\text{Bi}_{0.75-1.25}\text{Ag}_{0.03-0.15}$
 25 - $\text{Pd}_{1.25-1.75}\text{Bi}_{1.25-1.75}\text{Co}_{0.005-0.02}$
 - $\text{Pd}_{0.4-1.0}\text{Rh}_{1.0-1.75}\text{Bi}_{0.75-1.25}$
 - $\text{Pd}_{0.15-2.25}\text{Rh}_{0-2.5}\text{Bi}_{0.15-2.75}$

30 on a carrier, wherein the indices indicate the mass ratios of the respective elements.

7. The use of a catalytically active composition comprising an active component of the formula $\text{Pd}_a\text{Bi}_c\text{Y}_d$ according to claim 1 on a carrier for the dehydrogenation of cyclic or acyclic carbonyl compounds to the corresponding α,β -unsaturated carbonyl compounds.
 35

8. The use according to claim 7 wherein the cyclic or acyclic carbonyl compound is selected from the group consisting of cyclopentanone, butanone, butyraldehyde, cyclohexanone and isovaleraldehyde.
- 5 9. The use of a catalytically active composition comprising an active component of the formula $\text{Pd}_a\text{Rh}_b\text{Bi}_c$ according to claim 2 on a carrier for the dehydrogenation of cyclic or acyclic carbonyl compounds to the corresponding α,β -unsaturated carbonyl compounds.
- 10 10. The use according to claim 9, wherein the cyclic or acyclic carbonyl compound is selected from the group consisting of cyclopentanone, butanone, butyraldehyde, cyclohexanone and isovaleraldehyde.
- 15 11. The use of a catalytically active composition comprising an active component of the formula Pd_aBi_c according to claim 3 on a carrier for the dehydrogenation of cyclic or acyclic carbonyl compounds to the corresponding α,β -unsaturated carbonyl compounds.
- 20 12. The use according to claim 11, wherein the cyclic or acyclic carbonyl compounds are selected from the group consisting of cyclopentanone, butanone, butyraldehyde, cyclohexanone and isovaleraldehyde.
- 25 13. The use of a catalytically active composition comprising an active component of the formula $\text{Pd}_a\text{Rh}_b\text{Bi}_c\text{Z}_e$ according to claim 4 on a carrier for the dehydrogenation of cyclic or acyclic carbonyl compounds to the corresponding α,β -unsaturated carbonyl compounds.
- 30 14. The use according to claim 13, wherein the cyclic or acyclic carbonyl compounds are selected from the group consisting of cyclopentanone, butanone, butyraldehyde, cyclohexanone and isovaleraldehyde.
- 35 15. The use of a catalytically active composition comprising an active component of the formula $\text{Pd}_a\text{Bi}_c\text{Co}_e$ according to claim 5 on a carrier for the dehydrogenation of cyclic or acyclic carbonyl compounds to the corresponding α,β -unsaturated carbonyl compounds.

- 54 -

16. The use according to claim 15, wherein the cyclic or acyclic carbonyl compounds are selected from the group consisting of cyclopentanone, butanone, butyraldehyde, cyclohexanone and isovaleraldehyde.
- 5 17. The use of a catalytically active composition comprising an active component of the formula:
- 10
- $\text{Pd}_{0.5-1.0}\text{Rh}_{0.5-1.25}\text{Bi}_{1.25-1.75}\text{Ag}_{0.05-0.15}$
 - $\text{Pd}_{0.5-1.0}\text{Rh}_{1.0-1.5}\text{Bi}_{0.75-1.25}\text{Pt}_{0.01-0.1}$
 - $\text{Pd}_{0.25-0.5}\text{Rh}_{1.75-2.5}\text{Bi}_{0.25-0.5}\text{Co}_{0.01-0.1}$
 - $\text{Pd}_{0.5-1.25}\text{Rh}_{0.5-1.25}\text{Bi}_{0.75-1.5}\text{Cr}_{0.01-0.1}$
 - $\text{Pd}_{1.0-1.75}\text{Rh}_{0.25-0.75}\text{Bi}_{0.75-1.5}\text{Pt}_{0.0-0.15}\text{Co}_{0.01-0.1}$
 - $\text{Pd}_{1.0-1.75}\text{Rh}_{0.25-0.75}\text{Bi}_{0.75-1.5}\text{Pt}_{0.05-0.15}$
 - $\text{Pd}_{0.5-1.0}\text{Rh}_{1.0-1.75}\text{Bi}_{0.5-1.25}\text{Ag}_{0.03-0.15}\text{Ca}_{0.02-0.1}$

15

 - $\text{Pd}_{0.4-1.0}\text{Rh}_{1.0-1.75}\text{Bi}_{0.75-1.25}\text{Ag}_{0.03-0.15}$
 - $\text{Pd}_{1.25-1.75}\text{Bi}_{1.25-1.75}\text{Co}_{0.005-0.02}$
 - $\text{Pd}_{0.4-1.0}\text{Rh}_{1.0-1.75}\text{Bi}_{0.75-1.25}$
 - $\text{Pd}_{0.15-2.25}\text{Rh}_{0-2.5}\text{Bi}_{0.15-2.75}$

20

on a carrier for the dehydrogenation of cyclic or acyclic carbonyl compounds to the corresponding α,β -unsaturated carbonyl compounds.

18. The use according to claim 17, wherein the cyclic or acyclic carbonyl compounds are selected from the group consisting of cyclopentanone, butanone, butyraldehyde, cyclohexanone and isovaleraldehyde.

25